

Resolving Bulimia Nervosa Using An Innovative Neural Therapy Approach: Two Case Reports

Michael I Gurevich, MD

Private practice in, psychiatry and integrative medicine

997 Glen Cove Avenue, Glen Head, NY 11545

516-6749489 office, 516-7595946 fax

MIGurevich@gmail.com

Corresponding author

Myung Kyu Chung, MD

Clinical Professor of Family Medicine

Cooper Medical School of Rowan University

Chung Institute of Integrative Medicine

110 Marter Avenue, Suite 507

Moorestown, NJ 08057

(856)222-4766 office, (856)222-1137 fax

dockyu@gmail.com

Patrick J LaRiccia, MD, MSCE

Adjunct Scholar Center for Clinical Epidemiology and

Biostatistics at the Perlman School of Medicine of the

University of Pennsylvania

Research Director Won Sook Chung Foundation

Penn-Presbyterian Medical Center

51 N. 39th St.

Philadelphia, Pa 19104

lariccip@mail.med.upenn.edu

Running title: Resolving Bulimia with Neural Therapy

Ethical Approval: Our study does not require ethical approval.

Key Clinical Message

Conventional treatment of Bulimia Nervosa is long term, expensive, and often ineffective. Neural Therapy holds promise for treating Bulimia Nervosa in a shorter term, lower cost, and more effective manner. Much of Neural Therapy involves the superficial injection of local anesthetic injections. Implementation into current practice would be feasible.

Abstract

Objective: Conventional treatment of Bulimia Nervosa (BN) requires a multidisciplinary approach, hospitalizations, rehabilitation, and medications. Despite these interventions patients often relapse or discontinue therapy. Multiple psychiatric and medical complications are common. We explore a new application of Neural Therapy (NT) to treat Bulimia.

Methods: NT is an approach of treating multiple conditions using injections of local anesthetics superficially or deeper e.g. autonomic nervous system ganglia. Two case studies are reported.

Results: Herein for the first time are two case reports that illustrate rapid resolution of BN by including Neural Therapy (NT) in the treatment regime. In one case all BN symptoms resolved after just one session of injections; in the second case most symptoms resolved after the first NT treatment session while 4 additional treatment sessions achieved long lasting remission.

Conclusion: NT might be considered a promising therapeutic tool to treat BN. Further controlled research is indicated.

Key Words: Bulimia Nervosa, Neural Therapy, Local Anesthesia, Injections

Significant Outcomes: The symptoms of BN were rapidly resolved in two patients using Neural Therapy.

Limitations: This was a case study of two patients who had simultaneously received other treatments that may have confounded the results.

Introduction

Bulimia Nervosa (BN) is characterized by binge eating and compensatory behavior such as frequent self-induced vomiting¹. Lifetime prevalence is 1% worldwide², affecting women at 1.5% and men at 0.5%². BN is often accompanied by additional psychopathology, suicidal behavior³ and physical complications^{2,4}. Treatment for BN is complex: involving teams of practitioners⁵ including group, family⁶ and individual psychotherapists; dietary consultations; medical management of complications; psycho-pharmacotherapy; hospitalizations and day treatment programs^{7,8}. However treatment efficacy remains limited, with high dropout rates, frequent relapses and long lasting morbidity⁵⁻⁸.

We present two case reports of patients with BN who had failed conventional medical treatment. Their treatment regime included an innovative component called neural therapy (NT). This approach appears to have provided rapid recovery of cases in my practice (MG), and a possible new outlook on the pathophysiology of the condition.

Neural Therapy⁹ (NT) is a method that uses injections of local anesthetic into trigger points, scars, peripheral nerves, autonomic ganglia, tendon and ligament insertions, the epidural space and tissues. The determination of injection site(s) depends on the findings of a conventional medical evaluation in the context of referred pain, dermatomes, regional influence of autonomic ganglia and identified interference fields (IFs). IFs can be generated by any damaged tissue via chronic stimulation of afferent neurons in the autonomic nervous system resulting in chronic autonomic reflex activity such as nausea, vomiting, and pain. Scars are often sources of interference fields.

NT was developed in Germany and the former USSR in the first half of the 1900's. It is commonly used in German and Spanish speaking countries, but poorly known in the English speaking world. It has been used for the treatment of multiple medical conditions, injuries, and pain¹⁰⁻²¹. However, its application in the treatment of psychiatric conditions is largely

unexplored. The research from VA hospitals to treat veterans suffering from PTSD using Stellate Ganglion injections^{22,23} (which is a NT technique) shows very promising results. It has been used as an adjunct therapy for Bipolar Disorder patients with good results.^{24,25} A literature search in PubMed (which includes MEDLINE), EMBASE, AMED, and CINAHL found no prior report of the use of NT in the treatment of BN. Both patients mentioned herein have provided signed written consent for publication of this report.

Case presentations:

A 48-year-old married female executive presented with persistent daily vomiting for the last 20 years. She self-induced vomiting nightly. It has been her compulsive habit, which she has accepted to maintain stable weight, guilt over eating large portions at night and relief of uncomfortable sensations in her stomach. She had suffered from bulimia and anorexia since age 16. At that time, she required extensive therapy managing distorted body images, excessive and compulsive food consumption, and vomiting to control her weight. Over the last several years her weight had been stable on a highly restricted diet. But any significant bending forward caused her to vomit. She reported an inability to feel emotions and cry. She was sexually abused at age 11 and abused drugs and alcohol in her late teens. Her father committed suicide when she was 6 years old. He suffered from alcoholism and depression. She tried multiple conventional therapies but felt limited or no improvement. Despite her psychiatric issues she excelled academically and professionally.

Her past medical history included insomnia, depression, hypothyroidism, gastro-esophageal reflux and other digestive problems. She received multiple courses of antibiotics due to frequent infections and food poisonings. She ingested two glasses of wine nightly to cope with insomnia. She was taking medications: Escitalopram, Ambien, Levothyroxine, Famotidine, Omeprazole and Calcium Carbonate. Her initial DSM5 psychiatric diagnoses were: Anorexia Nervosa in full remission, Bulimia Nervosa, Moderate; PTSD; Substance Use Disorder in full remission; and Major Depression, Mild. Her Clinical Global Impression-Severity (CGI-S) score was 5.

Neural therapy injections using a total of 10 ml of preservative-free 1% procaine, were injected into the following locations: scar on the right side of the face -1 ml; the umbilical scar-2 ml; intra-cutaneous blebs over the area of the esophagus-2 ml; xyphoid process-1 ml; upper abdomen-2 ml; and periosteal injections into the sternum-2 ml. During injections, she was asked to focus on her emotional traumas, losses in her life and her compulsion to vomit resulting in strong emotional relief with a lot of tears.

Also, a modified form of EMDR (Eye Movement Desensitization and Reprocessing) was done to facilitate processing of emerging emotions. She was asked to follow with her eyes MG's hand making figure eights in the air, while focusing on stressful events resulting in further emotional release. That night she did not vomit.

The patient's follow-up appointment occurred two weeks later. She had discontinued Famotidine, Omeprazole and Calcium Carbonate, and later tapered off Escitalopram. There has not been a single episode of self-induced vomiting or vomiting associated with bending forward for the last 23 months of follow up. Her CGI-S score is 1. She has had at minimum monthly follow-up visits for other health concerns and prevention. Currently, she has weekly to bi-weekly appointments with an acupuncturist/naturopathic physician in MG's office.

Case #2

A 25-year-old white male presented with a chief complaint of five months of self-induced vomiting and 30-pound weight loss. Six months prior he had discontinued a gluten-free diet and began to indulge in rapidly consuming large amounts of junk food, and hiding this behavior from his family. He felt guilty and fearful of gaining weight and becoming fat. He resolved his fear and guilt by self-induced vomiting after each bingeing episode. The number of vomiting episodes quickly progressed to several times per day, becoming compulsive and unavoidable.

During vomiting episodes, he experienced emotional relief and almost sexual-like pleasure akin to a drug addiction. Initially, he felt gratified for controlling his weight during bingeing episodes but soon lost control, being unable to resist urges to vomit. This compulsion reminded him of the struggle he experienced earlier in his life coping with drug abuse. Compulsive behavior escalated over the preceding two months. He was anxious and preoccupied with the need to vomit, which he induced after each meal. He began to lose weight rapidly. His parents, concerned about unexplained weight loss, consulted a gastroenterologist. Gastroscopy revealed esophagitis. He was started on Omeprazole with some relief of heartburn and vomiting, but he developed severe constipation.

The patient had longstanding issues with anxiety and substance abuse. At age 23 he developed psychosis, depression and suicidal ideations, diagnosed by a consulting psychiatrist as schizophrenia. Several psychotropic medications were prescribed, but his parents refused medication treatment. MG was consulted, changing the diagnosis to drug-induced psychosis. He was treated using supplements, energetically based psychotherapy, dietary modifications, meditation and neural therapy. Keeping with the parents' desires, psychotropic medications were avoided. The patient had 27 office visits over a period of 28 months. He responded well but stopped follow-up a year and a half before the current visit. His DSM-5 diagnoses were Bulimia Nervosa, Extreme, Multiple Substance Use Disorder in full remission, Substance Induced Psychotic Disorder, in full remission, and Generalized Anxiety Disorder. His Clinical Global Impression-Severity (CGI-S) score was 6.

On this current visit, NT injections were applied to the skin overlying the stomach and esophagus ventrally and dorsally; the periosteum of the sternum; and the thyroid gland. 12 injections were done using 0.5 to 1 cc of preservative-free 1% procaine, in addition, 3 cc of preservative-free 1% procaine was given intravenously as an IV push of 1 to 3 minutes. The patient was also placed on: multivitamins, minerals, and omega 3 oils for general support; St. John's Wort and Avena Sativa for symptoms of depression; Soluna™ homeopathic-phytotherapy remedies # 3,8,17 and

20 for detoxification; and Biotonic™ Adaptogenic Tonic for adrenal support, to improve his digestion, decrease inflammation, and assist with anxiety and depression. On a visit one month later he reported cessation of bulimic episodes after the NT injections. His desire to compulsively vomit stopped. However, he could eat only small doses of food, experienced constipation, and indigestion, but was able to gain 2 lb. He continued Omeprazole for another month.

Neural therapy with localizations similar to the initial visit was repeated 4 times over the next two months in visits spaced 2-3 weeks apart. For the following nine months the patient was then seen monthly with the main therapeutic emphasis placed on persistent digestive issues, which resolved after a therapeutic trial of anti-parasitic medications, a gluten free diet and use of a mindful eating approach. In a total of 18 months of follow-up, the patient has been free of self-induced vomiting for 12 months. He had two episodes of self-induced vomiting at month six related to stress at work and persistent indigestion. At 9 months of follow up his CGI-S score was 1 and Clinical Global Impression-Improvement (CGI-I) score was 1.

Discussion:

Two adult cases of BN responded very quickly to outpatient therapy which included NT. A 48-year-old female responded after one NT treatment session and has been free of BN for the last 23 months. A 29-year-old male's self-induced vomiting was eliminated after the first visit. He had two vomiting episodes at the 6-month mark and has had no vomiting episodes the past 12 months.

The main limitations of this report are the small sample size, its retrospective nature with the possibility of selection bias, and the lack of controlled replication. Case reports sit on the bottom of evidence based medicine's hierarchy of study designs. One cannot generalize from such a small sample. Also, other treatments such as EMDR, food supplements and gluten restriction were administered, thus there is the problem of confounding which includes the possibility that

the whole treatment package rather than NT alone was responsible for the therapeutic benefit. Based on our clinical experience and training we feel that NT contributed the most to the positive outcomes. More vigorously designed future studies are needed to determine the contributions of the individual components and the entire regime as a whole.

Our total experience with BN includes seven cases. The other 5 cases were treated with psychotherapy, medications, and food supplements without success. Our positive experiences of NT after addition to our practices across all encountered medical problems prompts us to think that NT was a major contributor to the successful outcome of these two reported cases of BN.

BN treatment approaches encompass the application of behavioral modification, dietary changes and medication management for patients who are often reluctant to accept any extended therapy. Treatment strategies require long term management, efforts of multispecialty treatment teams, expensive hospitalizations and day treatment program management. High dropout rates are common, and full recovery remains problematic. There are no tactics that can offer speedy symptom resolution, are inexpensive and do not require long term compliance or patient commitment.

NT may offer a fresh understanding of BN's pathophysiological mechanisms and a highly effective treatment. It presents the possibility of a rapid resolution of compulsive vomiting and obsessive food behavior.

The pathophysiological mechanism of BN is complicated. A psychologically vulnerable population distressed by substance abuse, sexual and emotional traumas looks for relief from pain and suffering. Indulging in food offers a short-lasting pleasure. It is our hypothesis that self-induced vomiting coupled with food ingestion creates a Pavlovian conditioned response. After a number of repetitions, vomiting can become an independent provider of endorphin release, creating a vicious cycle. Part of this cycle is localized in the autonomic nervous system (ANS) within ganglions controlling the stomach and esophagus. Vulnerable tissue becomes part of an

interference field (IF). The IF in the damaged esophagus and stomach has a memory of its own: vomiting creates a pleasure experience. Thus a vicious cycle, akin to narcotic seeking behavior is formed and maintained by the IF.

The IF concept was developed to explain some of the of the NT effects. Scars, fractures, and dental lesions can create ANS disturbances that can affect other organs. Signals from them can provide non-linear and non-anatomically connected interference with affected organs. Over the past 90 years, several reports have documented clinical experience suggesting that injecting an IF such as a scar can resolve pain, organ dysfunction, or other symptoms near or at a distance from the IF.⁹⁻²⁵

Local anesthetic injections, like procaine have the capacity to re-polarize the nerve cell membrane resulting in restoration of organ function.⁹ Nerve cell repolarization is posited to be the crucial mechanism⁹ in contrast to endorphin release in acupuncture.

The rationale for using NT to treat BN is based on the hypothesis that BN is associated with an IF in autonomic nervous system ganglions affecting the esophagus and stomach. The ectodermal layer of the skin is connected to the ANS ganglions controlling those organs. By injecting the ectodermal layer of the skin the organ function becomes restored. Thus symptoms of compulsive vomiting are eliminated or significantly reduced. "Lightning reaction" a term developed by NT founders denotes an immediate healing reaction. It was observed in both presented cases.

Compulsive vomiting stopped immediately in both cases, however, in the second case there were two additional episodes of vomiting reported 6 months after the treatment.

NT can be convenient for psychiatric patients: it does not require conscious patient participation, examination of underlying psychodynamic mechanisms or long term medication use.

Interactions between physician and patient can be relatively brief, and frequent long term follow-up visits are not required. Materials used in NT are inexpensive. Training to learn superficial injections is relatively brief. The method is remarkably safe with very rare side effects. It can be

easily incorporated into any treatment program. In my (MG) psychiatric practice I have witnessed profound effects in treatment resistant patients.^{24,25}

Summary:

The effectiveness of conventional BN treatment has been problematic. The two reported cases of successfully treated BN patients point to the possibility of NT being an effective therapeutic agent in the treatment of BN. In both cases patients were able to rapidly halt vomiting. There were no reported adverse events. Future rigorously designed studies are indicated. We are open to collaboration in this endeavor.

Author Contributions:

Michael I Gurevich, MD was the treating physician and wrote the first draft.

Myung Kyu Chung, MD is a collaborating colleague, neural therapy practitioner and lecturer. He critically reviewed all aspects of the manuscript with special attention to: the description of the basic mechanism of action of NT in the introduction; the description of the neural therapy techniques used in the case presentations; and the proposed mechanisms of action described in the discussion section.

Patrick J LaRiccia, MD, MSCE was engaged in the development of the discussion section, the literature review directly related to NT, content development of the introduction, and editing and formatting the content of the case presentations.

Acknowledgement:

We extend our appreciation to Karl Kjer, PhD for his manuscript editing and formatting.

References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5). American Psychiatric Association, Arlington 2013
2. Hudson JI, Hiripi E, Pope HG JR, Kessler RC. The prevalence and correlates of eating disorders in the National Comorbidity Survey Replication. *Biol Psychiatry* 2007; 61:348-358.
3. Paul T, Schroeter K, Dahme B, Nutzinger DO. Self-injurious behavior in woman with eating disorders. *Am J Psychiatry* 2002; 159:408-411
4. Crow S, Swigart S, Medical assessment, In: *Assessment of Eating Disorders*, Mitchel, JE, Peterson, CB (Eds), Guilford Publications, New York 2005, p.120
5. McIntosh VVW, Jordan J, Builic CM. specialist supportive clinical management. In: *The Treatment of Eating Disorders: A Clinical handbook*, Grilo CM, Mitchel JE (Eds), The Guilford Press, New York 2010
6. Couturier J, Kimber M, Szatmari P. Efficacy of family-based treatment for adolescents with eating disorders: a systemic review and meta-analysis. *Int J Eating Disord* 2013; 46:3-11
7. Whittal ML, Agras, Gould RA. Bulimia Nervosa: A meta- analysis of psychosocial and pharmacological treatments. *Behav Ther* 1999; 30:117-135
8. Shapiro JR, Berkman ND, Brownley KA, et.al. Bulimia Nervosa treatment: a systemic review of randomized control trials. *Int J eat Disorder* 2007; 40:321-336
9. Dosch P, Dosch MP. *Manual of Neural Therapy According to Huneke*. New York, NY: Thieme; 2007.
10. Brobyn TL, Chung MK, LaRiccia PJ. Neural Therapy: An Overlooked Game Changer for Patients Suffering Chronic Pain? *Pain Relief*. 2015; 4(3):1-4.
<http://dx.doi.org/10.4172/2167-0846.1000184>

11. Weinschenk S. Neural therapy—A review of the therapeutic use of local anesthetics. *Acupuncture and Related Therapies*. 2012;1:5-9.
12. Hui F, Boyle E, Vayda E. A randomized controlled trial of a multifaceted integrated complementary-alternative therapy for chronic herpes zoster-related pain. *Alternative Medicine Review* 2012;17(1):57–68.
13. Chung MK, Chung D, LaRiccia PJ. Tonguepiercing and chronic abdominal pain with nausea and vomiting – two cases. *Explore (NY)*. 2015;11(1):59-62.
14. Chung, MK, LaRiccia PJ. Successful treatment of chronic nausea and vomiting related to body piercing and tattooing with integrative medicine interventions *Holist Nurs Pract*. 2015;29(1):33-6.
15. Mermoud J, Fischer L, Stabu L. Patient satisfaction of primary care for musculoskeletal diseases: A comparison between Neural Therapy and conventional medicine. *BMC Complementary and Alternative Medicine*. 2008;8:33.
16. Fischer L, Ludin SM, Puente de la Vega K, Sturzenegger M. Neuralgia of the Glossopharyngeal Nerve in a Patient with Posttonsillectomy Scarring: Recovery after Local Infiltration of Procaine—Case Report and Pathophysiologic Discussion. *Case Reports in Neurological Medicine* Volume 2015, Article ID 560546, 5 pages <http://dx.doi.org/10.1155/2015/560546>
17. Weinschenk S, Brocker K, Hotz L, Strowitzski T, Joos S. Successful therapy of vulvodynia with local anesthetics. A case report. *Forsch Komplementmed* 2013;20:138–143 Published online: April 15, 2013 DOI: 10.1159/000350023
18. Saha FJ, Komhard K, Langhorsta J. Successful Endoscopic Neural Therapy of a Patient with Chronic Pain Syndrome after Rectum Gunshot Injury. *Forsch Komplementmed* 2014;21:310–313 Published online: September 12, 2014 DOI: 10.1159/000367719

19. Chung MK, LaRiccia PJ. Successful Integrative Medicine Assessment and Treatment of Chronic Pain Associated With Breast Surgery: A Report of 3 Cases. *Holist Nurs Pract*. 2017;31(1):21-29.
20. Fischer L, Pfister M. Wirksamkeit der Neuraltherapie bei überwiesenen Patienten mit therapieresistenten chronischen Schmerzen. *Schweiz. Zschr. GanzheitsMedizin* 2007;19(1)30–35.
21. Egli S, Pfister M, Sabina M, Ludin SM, Puente de la Vega K, Busato A, Fischer L. Long-term results of therapeutic local anesthesia (neural therapy) in 280 referred refractory chronic pain patients. *BMC Complementary and Alternative Medicine*. 2015;15:200 DOI 10.1186/s12906-015-0735-z
22. Lipov G, Joshi JR, Sanders S, et.al. A unifying theory linking the prolonged efficacy of the stellate ganglion block for the treatment of chronic regional pain syndrome (CRPS), hot flashes, and posttraumatic stress disorder (PTSD). *Medical Hypotheses*. 2009; 72:657-661.
23. Hicky A, Handling S, Pevney E, et.al. Stellate ganglion block for PTSD. *Am J Psychiatry*. 2012 Jul; 169(7):760-760.
24. Gurevich MI, Robinson CL. An Individualized approach to treatment-resistant bipolar disorder: a case series. *Explore* 2016; 12:237-245.
25. Gurevich MI, Robinson CL. Medication-free alternatives for long-term maintenance of bipolar disorder: a case series. *Glob Adv Health Med*. 2015;4(2):53-60. doi:10.7453/gahmj.2014.064.